

Найти допустимые экстремали функционалов:

$$1) \mathcal{J}[y] = \int_{-1}^0 (12xy - y'^2) dx$$
$$y(-1) = 1, \quad y(0) = 0$$

$$3) \mathcal{J}[y] = \int_0^1 y y'^2 dx$$
$$y(0) = 1, \quad y(1) = \sqrt[3]{4}$$

$$5) \mathcal{J}[y] = \int_{-1}^1 (y'^2 - 2xy) dx$$
$$y(-1) = -1, \quad y(1) = 1$$

$$6) \mathcal{J}[y] = \int_{-1}^0 (y'^2 - 2xy) dx$$
$$y(-1) = 0, \quad y(0) = 2$$

Пример:

Задача 2:

$$y[y(x)] = \int_0^1 \sqrt{y(1+y'^2)} dx$$

$$y(0) = y(1) = \frac{1}{\sqrt{2}}$$

Искр. ур-е Эйлера:

$$-\frac{d}{dx} \left(\frac{1}{2\sqrt{y(1+y'^2)}} \cdot 2y'y' \right) + \frac{1+y'^2}{2\sqrt{y(1+y'^2)}} = 0$$

Проводим:

$$-(y''y + y'y') \sqrt{y(1+y'^2)} - \frac{yy'}{\sqrt{y(1+y'^2)}} (y' +$$

$$+ (y'^3 + 2yy'y')) + \frac{1+y'^2}{2\sqrt{y(1+y'^2)}} = 0$$

$$(y(1+y'^2))' \rightarrow$$

$$+ \frac{-2(y''y + y'^2) \cdot y(1+y'^2) - yy'^2 - yy'^2 - 2y^2y'^2y''}{2(y(1+y'^2))^{3/2}} +$$

$$* \frac{1+y'^2}{2\sqrt{y(1+y'^2)}} = 0$$

$$2(y''y^2 + y''y^2y'^2 + yy'^2 + yy'^2) - yy'^2 - yy'^2 - 2y^2y'^2y''$$

$$- y - yy'^2 - yy'^2 - yy'^2$$

$$- 2yy'^2$$

$$2y''y^2 - yy'^2 - y = 0$$

$$y(2y''y - y'^2 - 1) = 0$$

$$y = 0$$

~~xxxxxxxxxxxx~~

$$y' = p(y)$$

$$* \frac{dp}{dy} \cdot \frac{dy}{dx}$$

$$y'' = \frac{dp}{dy} \cdot \frac{dy}{dx}$$

$$y'' = p \frac{dp}{dy}$$

$$2p \frac{dp}{dy} \cdot y - p^2 - 1 = 0$$

$$2p \frac{dp}{dy} \cdot y = p^2 + 1$$

$$2p \frac{dp}{p^2+1} = \frac{dy}{y}; \quad \ln(p^2+1) = \ln y + \ln x$$
$$p^2+1 = cy$$

$$\Rightarrow p = \sqrt{cy-1}$$

$$y' = \sqrt{cy-1}$$

$$\frac{dy}{\sqrt{cy-1}} = dx$$

$$(cy-1)^{-\frac{1}{2}} dy = x + C_1$$

$$\frac{2}{c} \sqrt{cy-1} = x + C_1$$

$$\frac{4}{c^2} (cy-1) = (x+C_1)^2$$

$$\frac{4}{c} y = (x+C_1)^2 + \frac{4}{c^2}$$

$$y = \frac{c}{4} (x+C_1)^2 + \frac{1}{c}$$

$$y(0) = \frac{c}{4} C_1^2 + \frac{1}{c} = \frac{1}{\sqrt{2}}$$

$$y(1) = \frac{c}{4} (1+C_1)^2 + \frac{1}{c} = \frac{1}{\sqrt{2}}$$